

The University of North Carolina
at Greensboro

JACKSON LIBRARY



CQ
no.856

Gift of:
June Irene Bowers
COLLEGE COLLECTION

BOWERS, JUNE IRENE. On Rational Extensions of a Ring. (1971)
Directed by Dr. Robert L. Bernhardt. Pp. 33.

Let R be a ring, and let M_R denote that M is a right R -module. In this paper, the unique maximal rational extension of R_R , denoted by \bar{R} , is shown to be a ring whose operations preserve the module structure in $(\bar{R})_R$. It is then shown that if the right singular ideal is zero, the injective envelope of R_R is the maximal rational extension and therefore is a ring. It is further shown that this ring is right self-injective and is a von Neumann ring.

These results are applied to the $n \times n$ upper triangular matrix rings to compute their injective envelopes.

ON RATIONAL EXTENSIONS OF A RING


by

June Irene Bowers

A Thesis Submitted to
the Faculty of the Graduate School at
The University of North Carolina at Greensboro
in Partial Fulfillment
of the Requirements for the Degree
Master of Arts

Greensboro
June, 1971

Approved by


Thesis Adviser

APPROVAL SHEET

This thesis has been approved by the following committee of
the Faculty of the Graduate School at the University of North
Carolina at Greensboro.

Thesis
Adviser

Robert L. Dombardt

Oral Examination
Committee Members

Hughes B. Hayde, III

E.E. Posey

J.T. Hagerkelly

February 22, 1971
Date of Examination

ACKNOWLEDGMENT

The author wishes to express her gratitude to Dr. Robert L. Bernhardt for his patience and valuable assistance during the writing of this thesis.

TABLE OF CONTENTS

	Page
INTRODUCTION AND PRELIMINARY REMARKS.	v
CHAPTER I: THE INJECTIVE ENVELOPE CONSIDERED AS A RATIONAL EXTENSION.	1
Section 1: Rational and Essential Extensions	1
Section 2: When is the Injective Envelope a Ring?	8
Section 3: Utumi Quotient Rings and von Neumann Rings. .	18
CHAPTER II: THE INJECTIVE ENVELOPE OF THE $n \times n$ UPPER TRIANGULAR MATRIX RING	28
SUMMARY	32
BIBLIOGRAPHY.	33

384961

INTRODUCTION AND PRELIMINARY REMARKS

The purpose of this paper is to examine the rational extensions of a ring and to prove that under certain conditions the injective hull of a ring is a right self-injective, von Neumann ring. Furthermore, we shall show that the module structure of the injective hull induces the ring multiplication.

In the first section of Chapter I, we define a rational extension of a module over a ring. This concept is closely related to essential extensions of modules in that every rational extension of a module is an essential extension. Several theorems that characterize rational extensions are proved. Using these theorems, we show that when M_R is a right R -module with zero right singular submodule, every essential extension of M_R is a rational extension.

In section 2, the concept of maximal rational extension is introduced. We prove that any right R -module M_R has a maximal rational extension unique up to isomorphism, and we denote it by \bar{M} . For any ring R with unity, the maximal rational extension \bar{R} of R is shown to be a ring whose operations preserve the module structure in \bar{R} . We show, using theorems established in section 1, that if R is a ring with zero right singular ideal, then the injective hull of R_R is $(\bar{R})_R$ and therefore is a ring.

We further characterize this ring in section 3. After defining right quotient rings and von Neumann rings, we show for any ring R with zero right singular ideal that the injective hull E of R_R is the maximal right quotient ring of R . We then prove $\Lambda = \text{Hom}_R(E, E)$ is a right self-injective, von Neumann ring. We conclude the chapter by showing that when the right singular ideal of a ring is zero, the injective hull of R is ring isomorphic to Λ , and is therefore a right self-injective, von Neumann ring.

The results of Chapter I are used in Chapter II to show that the injective hull of the $n \times n$ upper triangular matrix ring over a division ring D is the full ring of $n \times n$ matrices over D .

Throughout this paper, the term "ring" will mean "ring with unity".

We define a right R -module as an abelian group $(M, +)$ together with a function $\mu : M \times R \rightarrow M$, where we let $\mu(a, x) = a\mu x$ such that:

- (a) $x\mu(a+b) = x\mu a + x\mu b$ for all $a, b \in R$ and $x \in M$;
- (b) $(x+y)\mu a = x\mu a + y\mu a$ for all $x, y \in M$ and $a \in R$;
- (c) $x\mu(ab) = (x\mu a)\mu b$ for all $a, b \in R$ and $x \in M$;
- (d) $x\mu 1 = x$ for all $x \in M$.

We shall denote the right R -module M by M_R . If N is a subgroup of M , then N is itself a candidate for a right R -module. We shall call a subgroup N of M a submodule of M_R whenever $x\mu a$ belongs to N for all $a \in R$ and $x \in N$. This will be denoted by $N_R \subseteq M_R$ or $M_R \supseteq N_R$.

If M_R and L_R are right R -modules, a function $f : M \rightarrow L$ is called a right R -homomorphism from M into L if in addition to being a group homomorphism it satisfies the following:
 $f(x\mu_1 a) = f(x)\mu_2 a$ for all $a \in R$ and $x \in M$, where μ_1 and μ_2 represent the module operations for M and L respectively. We shall denote the set of all right homomorphisms from a right R -module A_R into a right R -module B_R by $\text{Hom}_R(A, B)$.

A submodule N_R of M_R is said to be an essential submodule of M_R provided if for every submodule $0 \neq L_R$ of M_R , $N \cap L \neq 0$. In case N_R is an essential submodule of M_R , we say M_R is an essential extension of N_R and denote it by $M \nabla N$.

A module M_R is said to be R -injective provided for every exact sequence of R -modules $0 \rightarrow A \xrightarrow{f} B$, that is $f : A \rightarrow B$ is a one-to-one R -homomorphism, and for every $\phi \in \text{Hom}_R(A, M)$, there exists a $\psi \in \text{Hom}_R(B, M)$ such that $\phi = \psi \circ f$.

A result known as Baer's Lemma is used in this paper to check the injectivity of a module and is stated as follows:

Baer's Lemma. For a module Q_R , the following are equivalent:

- (1) Q_R is injective;
- (2) For every right ideal I of R and for every R -homomorphism $\phi : I \rightarrow Q$, there exists a $\phi' \in \text{Hom}_R(R, Q)$ such that $\phi'(a) = \phi(a)$ for all $a \in I$;
- (3) For every right ideal I of R and for every R -homomorphism $\phi : I \rightarrow Q$, there is a $q \in Q$ such that $\phi(a) = qa$ for all $a \in I$.

We shall call an injective **essential** extension of a module M_R the injective envelope or injective hull of M_R and denote it by $E(M)$. It is a well-known result that every module has an injective essential extension, and if E and E' are two injective essential extensions of M , then there exists an R -isomorphism $f: E \rightarrow E'$ such that $f(x) = x$ for every $x \in M$. Thus the injective envelope of a module is unique up to isomorphism.

CHAPTER I

THE INJECTIVE ENVELOPE CONSIDERED AS A RATIONAL EXTENSION

Section 1: Rational and Essential Extensions

1.1.1 DEFINITION. If $M_R \supseteq N_R$, then M_R is a rational extension of N_R , denoted by $M \nabla N$, in case for each submodule B_R of M_R such that $N_R \subseteq B_R$, $f \in \text{Hom}_R(B, M)$ satisfies $f(N) = 0$ if and only if $f = 0$.

1.1.2 LEMMA. If $M \nabla N$, then $M \nabla N$.

Proof: Let M_R and N_R be right R -modules such that $M \nabla N$. Let F_R be a submodule of M_R . Suppose $F \cap N = 0$. It suffices to show $F_R = 0$. Since $F \cap N = 0$, $B = F + N$ is a direct sum. Let $a \in B$. Then there exist $x \in F_R$ and $y \in N_R$ such that $a = x + y$. So if $\pi_F(a) = \pi_F(x+y) = x$, then $\pi_F \in \text{Hom}_R(B, M)$ and it is clear that $\pi_F(N) = 0$. Therefore, since $M \nabla N$, $\pi_F = 0$. This implies $F = 0$ since $F = \text{Im } \pi_F$.

1.1.3 LEMMA. $M \nabla N$ if and only if for each $x \in M$ and $0 \neq y \in M$, there exist $r \in R$ and $n \in Z$ such that $xr + xn \in N$ and $yr + yn \neq 0$.

Proof: (\Leftarrow) Let B_R be a submodule of M_R such that $M_R \supseteq B_R \supseteq N_R$ and f be an element of $\text{Hom}_R(B, M)$ such that $f(N) = 0$. We need to show $f = 0$; so assume $f \neq 0$. Then, there exists an $x \in B$ such that $y = f(x) \neq 0$. By hypothesis, there is an $r \in R$ and $n \in Z$ such that $xr + xn \in N$ and

$yr + yn \neq 0$. However, $0 = f(xr+xn) = f(xr) + f(xn) = f(x) \cdot r + f(x) \cdot n = yr + yn$. This is a contradiction. Therefore, $f = 0$.

(+) Assume $M \nabla N$. Let x and $0 \neq y$ be elements of M_R . Let $x_N = \{(r,n) \mid (r,n) \in R \times Z \text{ and } xr + nx \in N\}$, and $y_0 = \{(r,n) \mid (r,n) \in R \times Z \text{ and } yr + ny = 0\}$. It is sufficient to show that $x_N \not\subseteq y_0$; that is, there exists $(r,n) \in x_N$ with $(r,n) \notin y_0$. To this end, we shall show if $y_0 \supseteq x_N$, then $y = 0$, and thus arrive at a contradiction since $y \neq 0$.

Assume $y_0 \supseteq x_N$. Let B denote the submodule of M which consists of elements of the form $a + x(r,n)$ for some $a \in N$ and $(r,n) \in R \times Z$ where $x(r,n) = xr + xn$; thus, $B = N + xR + xZ$. Consider the relation f between B and M defined by $f(a+x(r,n)) = y(r,n)$. To show f is a function, assume $a + x(r,n) = a' + x(r',n')$. Then $x(r-r',n-n') = a - a' \in N$. So, $(r-r',n-n') \in x_N$. Since $y_0 \supseteq x_N$, $yr - yr' + yn - yn' = 0$. This implies $yr + yn = yr' + yn'$. But $f(a+x(r,n)) = yr + yn$ and $f(a'+x(r',n')) = yr' + yn'$. Hence, $f(a+x(r,n)) = f(a'+x(r',n'))$. Now we need to show f is an R -homomorphism. Let b_1 and b_2 be elements of B . There exist $a_1, a_2 \in N$, $r_1, r_2 \in R$, and $n_1, n_2 \in Z$ such that $b_1 = a_1 + x(r_1, n_1)$ and $b_2 = a_2 + x(r_2, n_2)$. One property of an R -homomorphism is satisfied since $f(b_1+b_2) = y(r_1+r_2, n_1+n_2) = y(r_1, n_1) + y(r_2, n_2) = f(b_1) + f(b_2)$. Now if $r \in R$, $f(b_1 r) = f([a_1 + x(r_1, n_1)] \cdot r) = f(a_1 r + x r_1 r + n_1 x r) = f(a_1 r) + f(x r_1 r) + f(n_1 x r) =$

$f(a_1 r + x(0,0)) + f(0 + x(r_1 r, 0)) + n_1 f(0 + x(r_1, 0)) = 0 + y(r_1 r, 0) + n_1 y(r, 0) = y r_1 r + n_1 y r = (y r_1 + n_1 y) r = y(r_1, n_1) r = f(b) r$. Therefore, $f \in \text{Hom}_R(B, M)$.

Let $a \in N$. Then $f(a) = f(a + x(0,0)) = 0$ and so $f(N) = 0$. Therefore, $f = 0$. This implies $y(r, n) = 0$ for all $(r, n) \in R \times Z$ and this means $y = 0$. So we know if $y_0 \supseteq x_N$, then $y = 0$. Since $y \neq 0$, $y_0 \not\supseteq x_N$ and there exists $(r, n) \in R \times Z$ such that $xr + xn \in N$ and $yr + ny \neq 0$. This proves the theorem.

1.1.4 LEMMA. If $M \nabla N$, then for each submodule F_R of M_R , $F \nabla (F \cap N)$.

Proof: Let $M \nabla N$ and $F_R \subseteq M_R$. Let $x \in F$ and $0 \neq y \in F$. Then there is an $r \in R$ and $n \in Z$ such that $xr + nx \in N$ and $yr + ny \neq 0$. But $xr + nx$ is also in F because $x \in F$. Therefore, $xr + nx \in F \cap N$ and $yr + ny \neq 0$. By 1.1.3, $F \nabla (F \cap N)$.

1.1.5 DEFINITION. For a module M_R , let $Z(M) = \{x \in M \mid (0:x) \text{ is an essential right ideal of } R\}$. We call $Z(M)$ the singular submodule of M_R .

The following lemma shows $Z(M)$ is a submodule of M_R , and in the case when $M_R = R_R$, $Z(M)$ is a two-sided ideal of R and is called the right singular ideal of R .

1.1.6 LEMMA. $Z(M)$ is a submodule of M_R and $Z(R)$ is a two-sided ideal of R .

Proof: Let M_R be a right R -module and let $x, y \in Z(M)$ and $r \in R$. We need to show $x - y \in Z(M)$ and $xr \in Z(M)$. First, we

show $x-y \in Z(M)$. Choose $k \in (0:x) \cap (0:y)$. Then $xk = 0 = yk$. Hence, $(x-y)k = 0$ and $k \in (0:x-y)$. Therefore, $(0:x-y) \supseteq (0:x) \cap (0:y)$. Let L be a non-zero right ideal of R . Then $((0:x) \cap (0:y)) \cap L = (0:x) \cap ((0:y) \cap L)$ which is not equal to zero since $(0:y) \cap L \neq 0$ and $R \nabla (0:x)$. Therefore, $(0:x) \cap (0:y)$ is essential in R . Since $(0:x-y) \supseteq (0:x) \cap (0:y)$, $(0:x-y)$ is essential in R and $(x-y) \in Z(M)$. To finish the first part of the proof, let $k \in [(0:x):r]$. Then $rk \in (0:x)$ and $x(rk) = (xr)k = 0$ which implies $k \in (0:xr)$. Hence, $(0:xr) \supseteq ((0:x):r)$. Now we shall show that $((0:x):r)$ is essential in R . Let K be a nonzero right ideal of R . Show $K \cap [(0:x):r] \neq 0$.

(Case 1): $r \cdot K = 0$. If $r \cdot k = 0$ for all $k \in K$, then $x(rk) = 0$ for all $k \in K$ which implies $0 \neq K \subset [(0:x):r]$.

(Case 2): $r \cdot K \neq 0$. Since rK is a right ideal of R , $(0:x) \cap rK \neq 0$. Therefore, there is a $0 \neq rk \in rK \cap (0:x)$. But $rk \in (0:x)$ implies $0 = x(rk)$ and that $k \in ((0:x):r)$. Clearly $0 \neq k \in [(0:x):r] \cap K$.

In either case, $((0:x):r)$ meets K nontrivially. Hence $((0:x):r)$ is essential in R . Since $(0:xr) \supseteq ((0:x):r)$, $(0:xr)$ is also essential in R . Therefore, $xr \in Z(M)$ and $Z(M)$ is a submodule of M_R .

By the preceding, $Z(R)$ is a right ideal of R . But if $x \in Z(R)$ and $a \in R$, $(0:ax) \supseteq (0:x)$ which implies $ax \in Z(R)$. Therefore, $Z(R)$ is a two-sided ideal of R .

1.1.7 LEMMA. Let $M_R \supseteq N_R$.

- (1) $Z(N) = Z(M) \cap N$, and if $M \nabla N$, $Z(M) \nabla Z(N)$;
- (2) if $M \nabla N$, then $Z(M) = 0$ if and only if $Z(N) = 0$.

Proof: (1) Let $x \in Z(N)$. Then $x \in N$ implies $x \in M$ which implies $x \in Z(M)$ since $R \nabla (0:x)$. Therefore, $x \in Z(M) \cap N$ and $Z(N) \subset Z(M) \cap N$. Let $x \in Z(M) \cap N$. Then $x \in N$ and $R \nabla (0:x)$ which implies $x \in Z(N)$. Therefore, $Z(M) \cap N \subset Z(N)$ and hence, $Z(N) = Z(M) \cap N$. Assume $M \nabla N$; we shall show $Z(M) \nabla Z(N)$. Let $0 \neq S_R$ be a submodule of M_R and $0 \neq K_R$ be a submodule of S_R . Then $K \cap N \neq 0$ and $0 \neq K \cap N = (K \cap S) \cap N = K \cap (S \cap N)$. Hence, $S \nabla (S \cap N)$ for each submodule S_R of M_R . So, $Z(M) \nabla (Z(M) \cap N)$ and since $Z(M) \cap N = Z(N)$, $Z(M) \nabla Z(N)$. Part (2) is immediate.

1.1.8 THEOREM. If $M_R \supseteq N_R$ and $Z(N) = 0$, then $M \nabla N$ if and only if $M \nabla N$.

Proof: (\leftarrow) 1.1.2

(\rightarrow) Assume $M \nabla N$ and $Z(N) = 0$. We will show $M \nabla N$ by using 1.1.3. By 1.1.7, $Z(N) = 0$ implies $Z(M) = 0$. Let x and $0 \neq y \in M_R$. Let $(N:x) = \{r \in R \mid xr \in N\}$ and $(0:y) = \{r \in R \mid yr = 0\}$. We know $(0:y)$ is not an essential ideal of R since $Z(M) = 0$ and $y \neq 0$. If we can show $(N:x)$ is essential in R , we will know $(0:y) \not\subseteq (N:x)$, and hence there will exist

an $r \in (N:x)$ such that $r \notin (0:y)$; that is, $xr \in N$ and $yr \neq 0$. Thus, the theorem will be proved. First, we need to show $(N:x)$ is a right ideal of R . Let $r_1, r_2 \in (N:x)$. Since M is a R -module, $x(r_1 - r_2) = xr_1 - xr_2$. But because N is a R -module, xr_1 and $xr_2 \in N$. Therefore, $xr_1 - xr_2 \in N$ and $r_1 - r_2 \in (N:x)$. The product $x(r_1 \cdot r_2) = (xr_1)r_2 \in N$ since $r_1 \in (N:x)$ implies $xr_1 \in N$. Therefore, $(N:x)$ is a subring of R . To show $(N:x)R \subseteq (N:x)$, let $r_1 \in (N:x)$ and $r_2 \in R$. By the previous argument, $x(r_1 \cdot r_2) = (xr_1)r_2 \in N$ and $r_1 \cdot r_2 \in (N:x)$. We now show $(N:x)$ is essential in R . Let $0 \neq s \in R$; then xsR is a submodule of M_R . If $xs = 0$, $s \in (N:x)$ and $0 \neq s \in sR \cap (N:x)$. If $xs \neq 0$, then $xsR \neq 0$ and $N \cap xsR \neq 0$. So there is an element a in R such that $0 \neq xsa \in N$. Therefore, $0 \neq sa \in (N:x)$ and since $sa \in sR$, $sR \cap (N:x) \neq 0$. In either case, $(N:x) \cap sR \neq 0$, and so $R \vee (N:x)$.

When $Z(R) = 0$, this relationship between rational and essential extensions is very useful. In particular, when $Z(R) = 0$, we know the injective envelope of R_R is also a rational extension.

1.1.9 LEMMA. If $F \nabla M$, and F_R is R -isomorphic to K_R under a R -isomorphism ϕ , then $K \nabla \phi(M)$.

Proof: Suppose $F \nabla M$ and ϕ is an onto, one-to-one R -homomorphism of F into K . Let B_R be a submodule of K_R such that $K_R \supseteq B_R \supseteq \phi(M)$. Suppose $f \in \text{Hom}_R(B, K)$ such that $f(\phi(M)) = 0$; we must show $f(B) = 0$. Since $B \supseteq \phi(M)$ and ϕ

is one-to-one, $\phi^{-1}(B) \supseteq M$. Let k be the restriction of $\phi^{-1}f\phi$ to $\phi^{-1}(B)$. Then $k \in \text{Hom}_R(\phi^{-1}(B), F)$ and $k(M) = 0$ since $k(M) = \phi^{-1}(f(\phi(M))) = \phi^{-1}(0) = 0$. However, $F \nabla M$ implies that $k = 0$ which implies $k(\phi^{-1}(B)) = 0$. But this means $(\phi^{-1}f\phi^{-1})(B) = \phi^{-1}(f(B)) = 0$, and since ϕ^{-1} is 1-1, that $f(B) = 0$. Therefore, $K \nabla \phi(M)$.

Section 2: When is the Injective Envelope a Ring?

1.2.1 DEFINITION. If $G_R \supseteq M_R$, then G_R is a maximal rational extension of M_R provided:

- (a) G_R is a rational extension of M_R ;
- (b) if $F_R \supseteq M_R$ and $F \nabla M$, then the identity map from M into M can be extended to a monomorphism of F into G ;
- (c) if $F_R \supseteq G_R$ and $F \nabla M$, then $F = G$.

One might ask if every right R -module has a maximal rational extension. The answer is yes, and the following theorem shows the existence of such a maximal rational extension which is unique up to isomorphism.

1.2.2 THEOREM. Let E be the injective hull of M_R for some right R -module M_R . Let $\Lambda = \text{Hom}_R(E, E)$ and $M^\Lambda = \{\lambda \in \Lambda \mid \lambda(M) = 0\}$. Then $\bar{M} = \cap \{\ker \lambda \mid \lambda \in M^\Lambda\}$ is a maximal rational extension of M_R , and \bar{M} contains each rational extension of M_R that is contained in E . Also, if G_R is any maximal rational extension of M_R , then the canonical injection of M_R into \bar{M} can be extended to a monomorphism of G onto \bar{M} .

Proof: The proof is divided into five parts which are collectively a proof of the theorem. Let M and \bar{M} be defined as stated in the theorem.

- (1) Show $\bar{M} \nabla M$. We know \bar{M} is an R -module since it is the intersection of the kernels of R -homomorphisms. It is clear that

$M_R \subseteq \bar{M}_R$. Let $B_R \subseteq \bar{M}_R$ such that $M_R \subseteq B_R \subseteq \bar{M}_R$. Let $f \in \text{Hom}_R(B, \bar{M})$ such that $f(M) = 0$. Now E is injective and $B \subseteq \bar{M} \subseteq E$. Consider the following diagram:

$$\begin{array}{ccccc}
 0 & \rightarrow & B_R & \xrightarrow{i} & E \\
 & & f \downarrow & \nearrow f' & \\
 & & \bar{M} & & \\
 & & i \downarrow & & \\
 & & E & &
 \end{array}$$

The injectivity of E implies the existence of $f' \in \text{Hom}_R(E, E)$ such that f' restricted to B_R is f . Since $f'(M) = f(M) = 0$, $f' \in M^\wedge$. Therefore, $\ker f' \supseteq \bar{M} \supseteq B$ and $f'(B) = 0$. So $f(B) = 0$, and $\bar{M} \nabla M$.

(2) Show \bar{M} contains each rational extension of M in E .

Let $F_R \subseteq E_R$ such that $F \nabla M$. Let $t \in M^\wedge$ and $K = \{x \in F \mid t(x) \in M\}$. Assume $t(F) \neq 0$. Since $E \nabla M$, $t(F) \cap M \neq 0$. Therefore, there exists an $x^* \in F$ such that $0 \neq y = t(x^*) \in t(F) \cap M$. Now $0 \neq x^* \in K$ and $0 \neq t(x^*) \in t(K)$. Let $N = t(F) \cap M$. Then $K = t^{-1}(N) \cap F$, and is hence a submodule of F_R . Observe now that $M \subseteq K + M \subseteq F$. Therefore t_0 , the restriction of t to $K + M$, is a function such that $t_0(K+M) \subseteq F$ because if $x \in K$, $t(x) \in M \subseteq F$. Since $F \nabla M$ and $t_0(M) = 0$, it follows that $t_0(K+M) = 0$. This implies $t(K) = t_0(K) = 0$. But, this is a contradiction to the fact that $0 \neq t(x^*) \in t(K)$. Hence, $t(F) = 0$ for all $t \in M^\wedge$. Therefore, by definition of \bar{M} , $F \subseteq \bar{M}$.

(3) Show \bar{M} satisfies condition (b) of definition 1.2.1. Let $H \nabla M$. Let θ_K loosely denote the canonical injection of any module K into another set containing K . Consider the following diagram:

$$\begin{array}{ccccc}
 0 & \longrightarrow & M & \xrightarrow{\theta_M} & H \\
 & & \theta_M \downarrow & & \searrow \phi \\
 & & \bar{M} & & \\
 & & \theta_{\bar{M}} \downarrow & & \\
 & & E & &
 \end{array}$$

Because E is injective, there exists $\phi \in \text{Hom}(H, E)$ such that $\phi(M) = M$. But $H \nabla M$ implies $H \nabla \bar{M}$. Therefore, $\ker \phi \cap M \subseteq \ker(\theta_{\bar{M}} \circ \theta_M) = 0$ since the injection maps are one-to-one. Because $\ker \phi \subseteq H$ and $H \nabla \bar{M}$, this implies $\ker \phi = \{0\}$. Thus ϕ is one-to-one and H is R -isomorphic to $\phi(H)$. By 1.1.9, $\phi(H) \nabla \phi(M) = M$. Thus by part (2) of this proof, $\phi(H) \subseteq \bar{M}$. Therefore, ϕ is a monomorphism of H into \bar{M} which extends the identity map on M .

(4) Show \bar{M} satisfies part (c) of the definition of maximal rational extension. Suppose $H \supseteq \bar{M}$ and $H \nabla M$. Then $H \nabla \bar{M}$ which implies $H \subseteq E$. Therefore, $H \subseteq \bar{M}$ and $H = \bar{M}$.

(5) If G_R is any maximal rational extension of M , we want to show the identity map of M into \bar{M} can be extended to a monomorphism of G onto \bar{M} . Let G_R be any maximal rational extension of M . We know the injective hull of G , denoted by

$E(G)$, is an injective essential extension of M . Therefore, there exists an isomorphism $f : E(G) \rightarrow E$ such that f is the identity on M . Let M' be defined for $E(G)$ as \bar{M} was for E . Since G is a rational extension of M and $G \subseteq E(G)$, $G \subseteq M'$ (parts(1), (2)). But by part 4, $M' = G$. Now $f(M') \subseteq \bar{M}$ since $f(M') \cap f(M) = M$ by 1.1.9. Also, $f^{-1}(\bar{M}) \subseteq M'$ by the same reasoning. Therefore, $f(M') \subseteq \bar{M} = ff^{-1}(\bar{M}) \subseteq f(M')$ and $f(M') = \bar{M}$. Therefore, f induces an isomorphism from $M' = G$ onto \bar{M} which is the identity on M .

1.2.3 LEMMA. Let R be a ring and E the injective hull of R_R . Let $\Lambda = \text{Hom}_R(E, E)$; then E is a left Λ -module.

Proof: Clearly Λ is a ring under composition. Define $\mu : \Lambda \times E \rightarrow E$ by $f \mu x = f(x)$. To show (E, μ) is a Λ -module, let f and g be elements of Λ and e_1 and e_2 be elements of E . One can easily check $f\mu(e_1 + e_2) = (f\mu e_1) + (f\mu e_2)$ and $(f+g) \mu e_1 = (f\mu e_1) + (g\mu e_1)$. Also, $(f \circ g) \mu e_1 = (f \circ g)(e_1) = f(g(e_1)) = f\mu(g(e_1)) = f\mu(g\mu e_1)$ and hence, (E, μ) is a Λ -module.

1.2.4 LEMMA. Let $Q = \text{Hom}_\Lambda(E, E)$ where R is a ring and E is the injective hull of R_R ; then E is a right Q -module. In fact, E is a (Λ, Q) -bimodule.

Proof: In this proof, we are assuming Q is a ring under composition that is opposite of the usual definition of composition of functions; that is, if $x \in E$ and $q, p \in Q$, then $(x)(q \circ p) = ((xq))p$. Let $x, y \in E$ and $p, q \in Q$. Define $\mu^* : E \times Q \rightarrow E$ by $x\mu^*q = xq$ where xq denotes x acted on by q . It is easy to show $(x+y)\mu^*q = x\mu^*q + y\mu^*q$ and $x\mu^*(q+p) = x\mu^*q + x\mu^*p$.

Also, $x\mu^*(q \circ p) = x(q \circ p) = ((xq))p = xq\mu^*p = (x\mu^*q)\mu^*p$. Therefore, (E, μ^*) is a Q -module. Let $f \in \Lambda$, $x \in E$, and $q \in Q$. Since q is a Λ -homomorphism, $(f\mu x)\mu^*q = (f\mu x)q = f(xq) = f\mu(x\mu^*q)$ and E is a (Λ, Q) -bimodule.

1.2.5 THEOREM. If R is a ring, E is the injective hull of R , and \bar{R} is the maximal rational extension of R , then \bar{R} is a ring whose operation $\cdot' : \bar{R} \times \bar{R} \rightarrow \bar{R}$ induces the module operation $\mu : \bar{R} \times R \rightarrow \bar{R}$ in \bar{R}_R ; in particular, R is a subring of \bar{R} .

Proof: Let R be a ring and E the injective hull of R . Let \bar{R} be defined as in 1.2.2 where $\Lambda = \text{Hom}_R(E, E)$. Let $Q = \text{Hom}_\Lambda(E, E)$. By 1.2.4, E is a (Λ, Q) -bimodule which implies for each $\lambda \in \Lambda$, $x \in E$, and $q \in Q$, $[\lambda(x)]q = \lambda[(x)q]$. If $R^\Lambda = \{\lambda \in \Lambda \mid \lambda(R) = 0\}$, then by 1.2.2, $\bar{R} = \{m \in E \mid R^\Lambda(m) = 0\} = \cap \{\ker \lambda \mid \lambda \in R^\Lambda\}$, and is the maximal rational extension of R_R . Observe since $1 \in R$, $R^\Lambda = \{\lambda \in \Lambda \mid \lambda(1) = 0\}$.

We first show Q is a right R -module. Define a function $\phi : R \rightarrow Q$ by $\phi(r) = \rho_r$ where $\rho_r(x) = xr$ for each $x \in E$ and module multiplication is understood. Clearly, $\rho_r \in Q$ for each $r \in R$. Let $r_1, r_2 \in R$ and $x \in E$. Then $(x)\rho_{r_1+r_2} = x(r_1+r_2) = xr_1 + xr_2 = (x)\rho_{r_1} + (x)\rho_{r_2}$ and hence, $\phi(r_1+r_2) = \phi(r_1) + \phi(r_2)$. Now $\phi(r_1 \cdot r_2) = \rho_{r_1 \cdot r_2}$ and $(x)\rho_{r_1 \cdot r_2} = (xr_1) \cdot r_2 = (x)\rho_{r_1} \circ \rho_{r_2}$ which implies $\phi(r_1 \cdot r_2) = \rho_{r_1} \cdot r_2 = \rho_{r_1} \circ \rho_{r_2} = \phi(r_1) \circ \phi(r_2)$. Also, if $r_1 \neq 0$ and $\rho_{r_1} = 0$, then

$\rho_{r_1}(1) = 1 \cdot r_1 = 0$, which is bad. So, ϕ is a ring monomorphism, and Q contains a subring $R' = \phi(R)$ which is ring isomorphic to R . Define $\mu' : Q \times R \rightarrow Q$ by $q\mu'r = q \circ \phi(r) = q \circ \rho_r$ for each $q \in Q$ and $r \in R$. The ring Q , together with μ' , is a R -module and moreover, Q is a ring whose multiplication (composition) induces the module multiplication, μ' .

Now we show Q_R is R -isomorphic to \bar{R}_R . If $q \in Q$, let $(1q) = q^\circ \in E$. Define $\theta : Q_R \rightarrow E_R$ by $\theta(q) = q^\circ$. If q_1 and q_2 are in Q , then $\theta(q_1 + q_2) = 1(q_1 + q_2) = 1q_1 + 1q_2 = \theta(q_1) + \theta(q_2)$. If $r \in R$, $\theta(q\mu'r) = (qr)^\circ = 1(qr) = 1(q \circ \phi(r)) = 1(q \circ \rho_r) = (1q)\rho_r = (q^\circ)\rho_r = q^\circ \cdot r = \theta(q) \cdot r$. Thus θ is a R -homomorphism. We want to show θ is an R -isomorphism. Let $q \in Q$ such that $\theta(q) = q^\circ = 0$ and let $x \in E$; define, for all $r \in R$, a function $k : R \rightarrow E$ by $k(r) = xr$ where module multiplication is understood. Since E is injective, there is a $\lambda \in \Lambda$ such that λ restricted to R is k . Now $\lambda(1) = k(1) = x$. So, $xq = (\lambda(1))q = \lambda(1q) = \lambda(q^\circ) = \lambda(0) = 0$. Therefore, since x was arbitrarily chosen in E , $q = 0$; hence $\ker \theta = \{0\}$ and θ is a monomorphism. Next, we want to show $\theta(Q) = \bar{R}$. Let $q \in Q$ and $\lambda \in \bar{R}^\wedge$. Then $q^\circ \in \theta(Q) = Q^\circ$ and $\lambda(q^\circ) = \lambda(1q) = (\lambda(1))q = (0)q = 0$. Since λ and q are arbitrary, $\bar{R}^\wedge(Q^\circ) = 0$ and $\theta(Q) = Q^\circ \subseteq \bar{R}$. Now we need to show $\bar{R} \subseteq Q^\circ$. Let $m \in E$. Define $\bar{\lambda}_m : R \rightarrow E$ by $\bar{\lambda}_m(r) = mr$ for each r in R . Since E is injective, there is a $\lambda_m \in \Lambda$ such that λ_m restricted to R

is $\bar{\lambda}_m$. Clearly, $\lambda_m(1) = m$ and, hence, for each m in E , there is a λ_m in Λ such that $\lambda_m(1) = m$. Fix $x \in \bar{R}$. Show $x \in Q^\circ$. Define $g : E \rightarrow E$ by $(m)g = \lambda_m(x)$ for each m in E . The function g is well-defined since if $\lambda'(1) = m$, then $(\lambda' - \lambda_m)(1) = 0$, which implies $\lambda' - \lambda_m \in R^\wedge$. Hence $(\lambda' - \lambda_m)(\bar{R}) = 0$ and then $(\lambda' - \lambda_m)(x) = 0$. So, $\lambda_m(x) = \lambda'(x)$ and g is well-defined. To show g is a Λ -homomorphism, let $\lambda' \in \Lambda$, $m \in E$ and show $(\lambda'(m))g = \lambda'(mg)$. By definition, $(\lambda'(m))g = \lambda_{\lambda'(m)}(x)$ and if $r \in R$, $\lambda_{\lambda'(m)}(r) = \lambda'(m) \cdot r = \lambda'(mr) = \lambda' \circ \lambda_m(r)$. So $(\lambda_{\lambda'(m)} - \lambda' \circ \lambda_m) \in R^\wedge$ which implies $\lambda_{\lambda'(m)}(y) = \lambda' \circ \lambda_m(y)$ for each $y \in \bar{R}$. Now $(\lambda'(m))g = \lambda_{\lambda'(m)}(x) = \lambda' \circ \lambda_m(x) = \lambda'((m)g)$, and $g \in Q$. Let i_E denote the identity on E ; then $(\lambda_1 - i_E)(R) = 0$ and so $(\lambda_1 - i_E)(\bar{R}) = 0$. This means $\lambda_1(x) = i(x) = x$. So $x = \lambda_1(x) = 1g = g^\circ \in Q^\circ$ and $\bar{R} \subseteq Q^\circ$. Therefore, $\bar{R} = Q^\circ$ and Q is R -isomorphic to \bar{R} .

One can show $(\bar{R}, +, \cdot')$ is a ring where $(r_1 + r_2) = \theta(\theta^{-1}(r_1) + \theta^{-1}(r_2))$ and $r_1 \cdot' r_2 = \theta(\theta^{-1}(r_1) \circ \theta^{-1}(r_2))$ (where θ is the R -isomorphism between Q and \bar{R}). So, \bar{R} is a ring whose ring multiplication induces the R -module multiplication, $\mu : \bar{R} \times R \rightarrow \bar{R}$, since if $r \in R$, then $\theta^{-1}(r) = \rho_r$ and $q^\circ \cdot' r = \theta(\theta^{-1}(q^\circ) \circ \theta^{-1}(r)) = \theta(q^\circ \rho_r) = \theta(q\mu'r) = \theta(q)\mu r = q^\circ \mu r$ for all $q^\circ \in \bar{R}$.

Let $"\cdot"$ denote multiplication in a ring R . The right regular module, R_R , is constructed by defining $r\mu r' = r \cdot r'$. Note that the module multiplication of the injective hull of R_R

extends μ , and since \bar{R}_R is a submodule of the injective hull, the module multiplication of \bar{R}_R extends μ and hence multiplication in R . So, if $r_1, r_2 \in R$, $r_1 \cdot' r_2 = r_1 \mu r_2 = r_1 \cdot r_2$ and R is a subring of \bar{R} .

1.2.6 THEOREM. Let R be a ring and E the injective hull of R_R . If $Z(R) = 0$, then E is a ring whose multiplication operation $\cdot' : E \times E \rightarrow E$ induces the module multiplication $\mu : E \times R \rightarrow E$.

Proof: Let R be a ring with $Z(R) = 0$. Let \bar{R} be defined as in 1.2.2 and 1.2.5. Since $\bar{R} \nabla R$ implies $\bar{R} \nabla R$, $\bar{R} \subseteq E$. However, $Z(R) = 0$ implies $E \nabla R$ which implies $E \subseteq \bar{R}$. Therefore, $\bar{R} = E$ and by 1.2.5, the theorem is proved.

The preceding theorems have proved that $Z(R) = 0$ is a sufficient condition for the injective hull to be a ring. The question "Is $Z(R) = 0$ a necessary condition" is answered in a paper by B. L. Osofsky [4]. In this paper, she gives an example of a ring with $Z(R) \neq 0$ whose injective hull is a ring. Thus, the answer to the above question is "no".

1.2.7 THEOREM. If R is a commutative ring, then \bar{R} is a commutative ring.

Proof: Let $(R, +, \cdot)$ be a commutative ring and $(\bar{R}, +, \cdot')$ be defined as in 1.2.5. By 1.2.6 we know R is a subring of \bar{R} . Fix $r \in R$. Define $\phi_1 : \bar{R} \rightarrow \bar{R}$ by $\phi_1(b) = r \cdot' b$ and $\phi_2 : \bar{R} \rightarrow \bar{R}$ by $\phi_2(b) = b \mu r$ for each $b \in \bar{R}$ where \cdot' induces μ . To check that ϕ_1 and ϕ_2 are R -homomorphisms, let a and

$b \in \bar{R}$ and $r_1 \in R$. Then $\phi_1(a+b) = r \cdot (a+b) = r \cdot a + r \cdot b = \phi_1(a) + \phi_1(b)$ and $\phi_2(a+b) = (a+b)\mu r = a\mu r + b\mu r = \phi_2(a) + \phi_2(b)$. Also, $\phi_1(a\mu r_1) = r \cdot (a\mu r_1) = r \cdot (a \cdot r_1) = (r \cdot a) \cdot r_1 = \phi_1(a) \mu r_1$, and $\phi_2(a\mu r_1) = (a\mu r_1) \mu r = a\mu(r_1 \cdot r) = a\mu(r \cdot r_1) = (a\mu r) \mu r_1 = \phi_2(a) \mu r_1$, since R is commutative. Therefore, ϕ_1 and $\phi_2 \in \text{Hom}_R(\bar{R}, \bar{R})$. Now let $z \in R$. Observe that $(\phi_1 - \phi_2)(z) = \phi_1(z) - \phi_2(z) = r \cdot z - z \mu r = r \cdot z - z \cdot r = r \cdot z - z \cdot r = 0$, since R is a subring of \bar{R} and R is commutative. Therefore, if $z \in R$, $(\phi_1 - \phi_2)(z) = 0$ and $(\phi_1 - \phi_2)(R) = 0$. However, $\bar{R} \nabla R$ implies $(\phi_1 - \phi_2)(b) = 0$ for each $b \in \bar{R}$ which implies $r \cdot b = b \mu r = b \cdot r$ for each $b \in \bar{R}$. Since r was arbitrarily chosen, we know $r \cdot b = b \cdot r$ for each $r \in R$, $b \in \bar{R}$; that is, elements of R commute with elements of \bar{R} .

Now we are ready to show \bar{R} is commutative. Fix $\bar{r} \in \bar{R}$. Define $f : \bar{R} \rightarrow \bar{R}$ by $f(b) = \bar{r} \cdot b$ for each b in \bar{R} and $g : \bar{R} \rightarrow \bar{R}$ by $g(b) = b \cdot \bar{r}$ for each $b \in \bar{R}$. We must show $f, g \in \text{Hom}_R(\bar{R}, \bar{R})$. The additive property clearly works. If $r \in R$, $f(b\mu r) = \bar{r} \cdot (b\mu r) = \bar{r} \cdot (b \cdot r) = (\bar{r} \cdot b) \cdot r = f(b)\mu r$ and $g(b\mu r) = (b\mu r) \cdot \bar{r} = (b \cdot r) \cdot \bar{r} = b \cdot (r \cdot \bar{r}) = b \cdot (\bar{r} \cdot r) = (b \cdot \bar{r}) \cdot r = g(b) \mu r$, since R commutes with \bar{R} . Thus, f and g are R -homomorphisms. Let $r \in R$. We see that $(f-g)(r) = \bar{r} \cdot r - r \cdot \bar{r} = \bar{r} \cdot r - \bar{r} \cdot r = 0$. Therefore, $(f-g)(R) = 0$, which implies (since $\bar{R} \nabla R$) that $(f-g)(\bar{R}) = 0$; that is, $\bar{r} \cdot b =$

$b \cdot \bar{r}$ for all $b \in \bar{R}$. Since \bar{r} was arbitrarily chosen, this is true for each $\bar{r} \in \bar{R}$. This means \bar{R} is commutative.

Section 3: Utumi Quotient Rings and von Neumann Rings

If R is a ring such that $Z(R) = 0$, in this section we will show that the injective hull of R_R is a maximal right quotient ring of R . We will also show that under this condition, the injective hull of R_R is a right self-injective, von Neumann ring.

1.3.1 DEFINITION. A ring A containing a ring R is a (Utumi) right quotient ring of R in case A_R is a rational extension of R_R .

1.3.2 DEFINITION. A right quotient ring A of R is maximal in case given any right quotient ring T of R , there exists a ring monomorphism of T into A which induces the identity map on R .

In the proof of 1.2.5, R is shown to be a subring of \bar{R} and we already know \bar{R} is a rational extension of R ; hence, \bar{R} is a right quotient ring of R . Therefore, every ring has a right quotient ring.

1.3.3 LEMMA. If A and B are right quotient rings of R and if $\phi : A_R \rightarrow B_R$ is a module monomorphism which induces the identity map on R , then ϕ is a ring monomorphism of A into B .

Proof: Let A and B be right quotient rings of R and ϕ be a module monomorphism of A_R into B_R which induces the identity map on R . Let " \circ " denote the product in A and " \cdot " denote the product in B . If $a \in A$ and $r \in R$, then $\phi(a \circ r) =$

$\phi(a) \cdot r$. Since ϕ is a module monomorphism, consider A to actually be a submodule of B ; that is, identify a and $\phi(a)$. Then $a \circ r = a \cdot r$ for all $a \in A, r \in R$. Let $a \in A$. Define $f_1 : A_R \rightarrow A_R$ by $f_1(x) = a \circ x$ for all $x \in A$ and $f_2 : A_R \rightarrow B_R$ by $f_2(x) = a \cdot x$ for all $x \in A$. Clearly, f_1 and f_2 are well-defined R -homomorphisms and so $f = f_1 - f_2$ is. Let $r \in R$. Then $f(r) = (f_1 - f_2)(r) = a \circ r - a \cdot r = 0$ and hence $f(R) = 0$. However, $B \nabla R$ then implies $f(A) = 0$ which implies $f(a_1) = a \circ a_1 - a \cdot a_1 = 0$ for all a_1 in A . Therefore, $a \circ a_1 = a \cdot a_1$ for all a_1 in A and since a was chosen arbitrarily, $a \circ a_1 = a \cdot a_1$ for all a and a_1 in A . Hence, $\phi(a \circ b) = a \circ b = a \cdot b = \phi(a) \cdot \phi(b)$ for all a and b in A , which implies ϕ is a ring monomorphism.

1.3.4 THEOREM. If A is a right quotient ring of R , there exists a ring monomorphism ϕ of A into \bar{R} which induces the identity map on R .

Proof: Let A be a right quotient ring of R . Then since $A \nabla R, A \nabla R$ and therefore $A \subseteq E(R)$, the injective hull of R . By 1.2.2, \bar{R} is a maximal rational extension of R in $E(R)$. Thus $i : R \rightarrow R$ can be extended to a module monomorphism $\phi : A_R \rightarrow \bar{R}_R$. But, by 1.3.3, ϕ is a ring monomorphism.

The preceding theorem tells us if R is a ring, \bar{R} is a maximal right quotient ring of R . So if $Z(R) = 0, E(R)$ is a maximal right quotient ring of R by 1.2.6.

1.3.5 COROLLARY. If R is a ring, then any two maximal right quotient rings of R are ring isomorphic under a mapping that induces the identity mapping on R .

Proof: Let A and B be maximal right quotient rings of a ring R . Then by definition of maximal right quotient ring, there exist ring monomorphisms ϕ_1 and ϕ_2 such that $\phi_1 : A \rightarrow B$ and $\phi_2 : B \rightarrow A$ which induce the identity map on R . Let i_A and i_B denote the identity maps on A and B , respectively. Now $(\phi_2 \circ \phi_1 - i_A)(R) = 0$, which implies, since $A \nabla R$, that $(\phi_2 \circ \phi_1 - i_A)(A) = 0$ which implies $\phi_2 \circ \phi_1 = i_A$. Likewise, $\phi_1 \circ \phi_2 = i_B$. Hence ϕ_1 is a ring isomorphism.

1.3.6 DEFINITION. A ring R is said to be von Neumann provided for each $a \in R$, there exists $x \in R$ such that $axa = a$.

1.3.7 DEFINITION. Let R be a ring. The Jacobson Radical of R , denote by $J(R)$, is the intersection of all the maximal right ideals of R .

1.3.8 DEFINITION. An element x of a ring R is said to be right quasi-regular provided there exists an element $x' \in R$ such that $x + x' + xx' = 0$ or equivalently, $(1+x)(1+x') = 1$.

The definitions of left quasi-regular and quasi-regular are analogous. A two-sided ideal I of R is said to be left quasi-regular provided every element of I is left quasi-regular. It is well-known that if R is a ring, $J(R)$ is a left (and right) quasi-regular two-sided ideal containing each left quasi-regular

two-sided ideal of R . The proofs will be omitted here but may be found in [3].

1.3.9 DEFINITION. A module M_R is said to be quasi-injective in case each homomorphism of any submodule of M_R into M_R can be extended to a homomorphism of M_R into M_R .

The following lemmas are proved in order to show the endomorphism ring, Λ , of a quasi-injective module is a right self-injective, von Neumann ring. We will use this theorem to prove that under certain conditions the injective hull of a ring is von Neumann and right self-injective.

1.3.10 LEMMA. A ring R is von Neumann if and only if for each $a \in R$, there exists an $e^2 = e \in R$ such that $aR = eR$.

Proof: (\rightarrow) Let R be a von Neumann ring. Choose $a \in R$. Then there exists $x \in R$ such that $axa = a$. Then $e = ax$ satisfies $e^2 = e$ and $eR = axR \subseteq aR$. But $a = ea \in eR$ so that $aR \subseteq eR$ and hence, $aR = eR$, where e is an idempotent.

(\leftarrow) Suppose $aR = eR$ for all $a \in R$, where e is an idempotent and depends on a . Let $a \in R$. Then there is $e^2 = e \in R$ such that $aR = eR$. So, there exists $r \in R$ such that $a = er$ and there exists $r' \in R$ such that $e = ar'$. Let $x = r'$. Then $axa = (a)(r')(er) = (ar')(er) = (e)(er) = e^2r = er = a$. Hence, R is von Neumann.

1.3.11 LEMMA. Let S be a von Neumann ring and let $P_r(S)$ denote the set of principal right ideals of S . Then

(1) if A and B are elements of $P_r(S)$, $A + B \in P_r(S)$;

(2) $P_r(S)$ contains each finitely generated right ideal of S .

Proof: (1) Let A and B be in $P_r(S)$. By 1.3.10, there exists $e = e^2$ in S and $f = f^2$ in S such that $A = eS$ and $B = fS$. Let $B_1 = (1-e)fS$. Now $A + B = \{eu + fv \mid u, v \in S\}$ and $A + B_1 = \{e(u' - fv) + fv \mid u', v \in S\}$. Choose $x \in A + B$. Then $x = eu + fv = eu + efv - efv + fv = e(u + fv - fv) + fv$, which implies $x \in A + B_1$. If $x \in A + B_1$, then $x = eu' + fv - efv = e(u' - fv) + fv$, which implies $x \in A + B$. Now, $B_1 = f_1S$ where $f_1 = (f - ef) = (1 - e)f$, and $f_1^2 = f_1 \in S$ and since $f_1 \in B_1$, $ef_1 = e(1 - e)f = (e - e^2)f = 0$. Now let $f' = f_1(1 - e)$. Then $f' \cdot f_1 = f_1(1 - e)f_1 = f_1f_1 - f_1ef_1 = f_1^2 = f_1$ and hence $f' \cdot f_1 = f_1$. So $(f')^2 = f'(f_1(1 - e)) = f_1(1 - e) = f'$. Since $f' = f_1(1 - e) \in f_1S$ and $f_1 = f'f_1 \in f'S$, it follows that $f'(S) = f_1(S) = B_1$. Hence, $A + B = eS + f'S$. Because $ef' = 0$ and $f'e = 0$, $A + B = eS + f'S = (e + f')(S) \in P_r(S)$.

(2) Suppose K is a finitely generated right ideal of S . Then there exist a_1, a_2, \dots, a_n in S such that $K = a_1S + \dots + a_nS$. But, 1.3.10 implies $a_iS = e_iS$, where e_i is an idempotent of S . Part (1) of this proof can be extended to finite sums. Hence K is a principal right ideal and is generated by an idempotent.

1.3.12 THEOREM. Let M_R be a quasi-injective module, $\Lambda = \text{Hom}_R(M, M)$, and $J = J(\Lambda)$. Then

(1) $J = \{\lambda \in \Lambda \mid M \cap \ker \lambda\}$ and Λ/J is von Neumann;

(2) if $J(\Lambda) = 0$, then Λ is a right self-injective, von Neumann ring.

Proof: Let $I = \{\lambda \in \Lambda \mid M \nabla \ker \lambda\}$. Let $\lambda \in \Lambda$ and $\alpha, \mu \in I$. One can easily show $\ker(\mu - \alpha) \supseteq \ker \mu \cap \ker \alpha$. Since $M \nabla (\ker \mu \cap \ker \alpha)$, $M \nabla \ker(\mu - \alpha)$, and thus $\mu - \alpha \in I$. Also, $\ker(\lambda \circ \alpha) \supseteq \ker \alpha$, which implies $M \nabla (\ker \lambda \circ \alpha)$. Hence, $\lambda \circ \alpha \in I$ and I is a left ideal of Λ . Let $0 \neq x \in M$. We wish to show $xR \cap \ker(\alpha \circ \lambda) \neq 0$. If $\lambda(xR) \neq 0$, then, because $\lambda(xR)$ is a submodule of M , $\lambda(xR) \cap \ker \alpha \neq 0$ since $M \nabla \ker \alpha$. Therefore, there exists $0 \neq xr \in xR$ such that $\lambda(xr) \neq 0$ and $\alpha(\lambda(xr)) = 0$. However, this means $0 \neq xr$ is in $\ker(\alpha \circ \lambda) \cap xR$. If $\lambda(xR) = 0$, then $\alpha(\lambda(xR)) = 0$. This implies $0 \neq xR \subseteq \ker(\alpha \circ \lambda)$ and so $xR \cap \ker(\alpha \circ \lambda) \neq 0$. In either case, $M \nabla \ker(\alpha \circ \lambda)$. Thus $\alpha \circ \lambda \in I$ and we can conclude that I is a two-sided ideal of Λ .

Let $\lambda \in I$. If $x \in \ker \lambda \cap \ker(1 + \lambda)$, then $\lambda(x) = 0$ and $x + \lambda(x) = 0$, which implies $x = 0$. Since $M \nabla \ker \lambda$, $\ker(1 + \lambda) = 0$. Thus we know for each $\lambda \in I$, $1 + \lambda$ is a one-to-one function and has a left inverse. Therefore, I is a left quasi-regular, two-sided ideal of R and consequently is a subset of J .

Now let $\lambda \in \Lambda$ and L_R be a submodule of M_R , maximal with respect to $N \cap \ker \lambda = 0$, where N is a submodule of M_R ; let $K = \ker \lambda$. Consider the relation ϕ , mapping $\lambda(L)$ into L , defined by $\phi(\lambda(x)) = x$ for each $\lambda(x) \in \lambda(L)$. If $x, y \in L$

such that $\lambda(x) = \lambda(y)$, then $\lambda(x-y) = 0$ and $x-y \in L \cap K = 0$. So $x = y$ and ϕ is a function. Since M is quasi-injective and $\lambda(L)_R \subseteq M_R$, ϕ is induced by some $\theta \in \Lambda$. Let $x + y = u \in L + K$ where $x \in L$ and $y \in K$. It is clear that $L + K \subseteq \ker(\lambda - \lambda\theta\lambda)$ since $(\lambda - \lambda\theta\lambda)(u) = \lambda(u) - \lambda\theta\lambda(u) = \lambda(x) - \lambda(x) = 0$.

Claim: $M \nabla (L+K)$.

Subproof: Assume $L + K$ is not essential in M . There exists a $N_R \subseteq M_R$ such that $(L+K) \cap N = 0$. We claim now that $(L+N) \cap K = 0$. Let $x \in L$ and $y \in N$ such that $0 \neq x+y \in (L+N) \cap K$. Then $(x+y) - x = y \in N \cap (L+K) = 0$. Therefore, $0 \neq x+y = x \in K \cap L = 0$, which is a contradiction since L is maximal with respect to missing K . Hence, $M \nabla (L+K)$.

Therefore $\lambda - \lambda\theta\lambda \in I$, which means for all $\lambda + I \in \Lambda/I$, there exists $\theta \in \Lambda$ such that $\lambda + I = \lambda\theta\lambda + I$. Hence, Λ/I is a von Neumann ring.

To finish part (1), it suffices to show $J \subseteq I$. Let $\lambda \in J$. By the previous argument, there exists a $\theta \in \Lambda$ such that $\lambda - \lambda\theta\lambda \in I$. Since J is left quasi-regular and $-\lambda\theta \in J$, there exists a $y \in \Lambda$ such that $y \circ (1 + \lambda\theta) = 1$. But $\lambda = y \circ (1 - \lambda\theta) \circ \lambda = y \circ (\lambda - \lambda\theta\lambda) \in I$, since I is an ideal. Therefore, $J \subseteq I$ and thus $J = I$ and (1) is proved.

(2) It follows from part (1) if $J = 0$, then $\Lambda/J = \Lambda/0 = \Lambda$ is a von Neumann ring. Lemma 1.3.11 will be used in the remainder of the proof to show Λ is right self-injective.

Let f be a Λ -homomorphism from a right ideal I of Λ into Λ . Define IM to be the submodule of M_R generated by $\{\lambda(m) \mid \lambda \in I \text{ and } m \in M\}$. If $x \in IM$, then there exist $n \in \mathbb{Z}^+, \lambda_1 \cdot \dots \cdot \lambda_n \in I$ and $m_1 \cdot \dots \cdot m_n \in M$ such that $x = \sum_{i=1}^n \lambda_i(m_i)$. Consider the function θ between IM and M defined by $\theta(x) = \sum_{i=1}^n f(\lambda_i)(m_i)$. We need to show θ is indeed a function. Suppose $y \in IM$. Then $y = \sum_{j=1}^t \mu_j(m_j)$, where $\mu_j \in I$ and $m_j \in M$ for $1 \leq j \leq t$. The right ideal of Λ generated by $\{\lambda_1, \lambda_2, \dots, \lambda_n, \mu_1, \mu_2, \dots, \mu_t\}$ is of the form $e\Lambda$ where $e^2 = e \in \Lambda$. Observe since $\lambda_1 \in e\Lambda$, $\lambda_1 = e\lambda$ for some $\lambda \in \Lambda$ and $e\lambda_1 \in e\Lambda$. Also, $e\lambda_1 = e(e\lambda) = e\lambda = \lambda_1$. Likewise $e\mu_j = \mu_j$, $f(\lambda_i) = f(e)\lambda_i$, and $f(\mu_j) = f(e)\mu_j$. Consequently, $\theta(x) = \sum_{i=1}^n f(\lambda_i)(m_i) = \sum_{i=1}^n (f(e)\lambda_i)(m_i) = f(e) \sum_{i=1}^n \lambda_i(m_i) = f(e) \cdot x$. Similarly, $\theta(y) = \sum_{j=1}^t f(\mu_j)(m_j) = f(e) \cdot y$. Thus if $x = y$, then $\theta(x) = f(e) \cdot x = f(e) \cdot y = \theta(y)$ and θ is a function. Clearly, θ is an R -homomorphism from IM into M . Since M is quasi-injective, θ is induced by some element $\theta' \in \Lambda$. Then $(\theta' \circ \lambda)(m) = \theta'(\lambda(m)) = f(\lambda)(m)$ for all $m \in M$ and $\lambda \in I$, so that $f(\lambda) = \theta' \circ \lambda$ for all $\lambda \in I$. This establishes that Λ satisfies Baer's condition for injectivity, and we conclude Λ is self-injective as a right Λ -module.

1.3.13 LEMMA. If R is a ring such that $Z(R) = 0$ and if E is the injective hull of R , then $\Lambda = \text{Hom}_R(E, E)$ is a right self-injective, von Neumann ring.

Proof: Let R be a ring such that $Z(R) = 0$. Let E denote the injective hull of R_R and $\Lambda = \text{Hom}_R(E, E)$. Let $\lambda \in J(\Lambda)$.

By 1.3.12, since injective implies quasi-injective, $E \nabla \ker \lambda$. But $\lambda(\ker \lambda) = 0$ which implies, since $E \nabla \ker \lambda$, that $\lambda(E) = 0$ or that $\lambda = 0$. Hence, $J(V) = 0$ and by 1.3.12, Λ is a right self-injective, von Neumann ring.

1.3.14 THEOREM. If R is a ring such that $Z(R) = 0$ and if E is the injective hull of R , then $\Lambda = \text{Hom}_R(E, E)$ is ring isomorphic to E .

Proof: Let R be a ring such that $Z(R) = 0$. Let E denote the injective hull of R_R and $\Lambda = \text{Hom}_R(E, E)$. We first want to show Λ is a right R -module. Let $r \in R$. Define $\eta_r : R_R \rightarrow E_R$ by $\eta_r(x) = r \cdot x$ for all $x \in R$, where " \cdot " is multiplication in E . Since E_R is injective and η_r is a R -homomorphism, η_r is induced by some $\eta_r^* \in \Lambda$. Define $\phi : R \rightarrow \Lambda$ by $\phi(r) = \eta_r^*$. Let $a, b \in R$. One can show without much difficulty that $[\eta_{a+b}^* - (\eta_a^* + \eta_b^*)](R) = 0$ and $(\eta_{a \cdot b}^* - \eta_a^* \circ \eta_b^*)(R) = 0$. Because $Z(R) = 0$, $E \nabla R$ and therefore $[\eta_{a+b}^* - (\eta_a^* + \eta_b^*)](E) = 0$ and $[\eta_{a \cdot b}^* - \eta_a^* \circ \eta_b^*](E) = 0$. Hence $\eta_{a+b}^* = \eta_a^* + \eta_b^*$ and $\eta_{a \cdot b}^* = \eta_a^* \circ \eta_b^*$ and ϕ is a ring homomorphism. Assume $r \in \ker \phi$; then $\eta_r^*(x) = 0$ for all $x \in E$. However, $1 \in R \subseteq E$; so, $\eta_r^*(1) = r = r \cdot 1 = 0$. Thus we know $\ker \phi = \{0\}$. Hence, Λ contains a subring isomorphic to R , namely $\phi(R)$. The ring Λ is a natural right R -module under the operation μ , where $\lambda \mu r = \lambda \circ \eta_r^*$ for all $\lambda \in \Lambda$ and $r \in R$. As in the proof of 1.2.5, for each $x \in E$, there is a $\eta_x^* \in \Lambda$ such that $\eta_x^*(1) = x$. Define a function $\theta : E \rightarrow \Lambda$ by $\theta(x) = \eta_x^*$. Clearly

θ is well-defined and $\theta(x+y) = \eta_{x+y}^* = \eta_x^* + \eta_y^*$ since
 $[\eta_{x+y}^* - (\eta_x^* + \eta_y^*)](r) = (x+y)r - xr - yr = 0$ for every $r \in R$
 and $E \nabla R$. It is not hard to see that $\eta_{x \cdot r}^* - \eta_x^* \mu r \in \text{Hom}_R(E, E)$
 and that $(\eta_{x \cdot r}^* - \eta_x^* \mu r)(1) = 0$. Thus, $(\eta_{x \cdot r}^* - \eta_x^* \mu r)(R) = 0$, which
 implies $(\eta_{x \cdot r}^* - \eta_x^* \mu r)(E) = 0$. So, $\theta(x \cdot r) = \eta_{x \cdot r}^* = \eta_x^* \mu r =$
 $\theta(x) \mu r$ and θ is an R -homomorphism. If $\theta(x) = \theta(y)$, then
 $x = \eta_x^*(1) = \eta_y^*(1) = y$ and θ is one-to-one. Choose $\lambda \in \Lambda$.
 Let $y = \lambda(1)$. Then $(\eta_y^* - \lambda) \in \text{Hom}_R(E, E)$, and $(\eta_y^* - \lambda)(1) = 0$,
 which implies $(\eta_y^* - \lambda)(R) = 0$. But again, $E \nabla R$ implies
 $(\eta_y^* - \lambda)(E) = 0$, which means $\theta(y) = \eta_y^* = \lambda$. Hence θ is a
 one-to-one, onto module homomorphism between E and Λ . So, Λ
 is a rational extension of R , where we identify R with $\phi(R)$,
 and Λ is thus a right quotient ring of R . Since θ is the
 identity on R , by 1.3.3 θ is a ring isomorphism and we can
 conclude E and Λ are ring isomorphic.

1.3.15 THEOREM. Let R be a ring such that $Z(R) = 0$.

Then the injective hull of R_R is a right self-injective, von Neumann ring which is also the maximal right quotient ring of R .

Proof: Let R be a ring such that $Z(R) = 0$. Let E denote the injective hull of R_R . By 1.2.6, $E = \bar{R}$ is a ring that contains R as a subring and is the maximal rational extension of R_R , hence, by 1.3.4 the maximal right quotient ring of R . By 1.3.13, Λ is a right self-injective, von Neumann ring and so, by 1.3.14, E is also.

CHAPTER II

THE INJECTIVE ENVELOPE OF THE $n \times n$ UPPER TRIANGULAR
MATRIX RING

This chapter is devoted to showing the injective hull of the $n \times n$ upper triangular matrix ring is the full $n \times n$ matrix ring.

In the following proofs, let D represent a division ring, $M_n(D) = S$ denote the full $n \times n$ matrix ring over D , and $UT_n(D) = K$ be the $n \times n$ upper triangular matrix ring over D . The matrix with 1 in the i^{th} row and j^{th} column with zero everywhere else will be denoted by E_{ij} . Also, $E(K)$ will denote the injective hull of K .

2.1 LEMMA. $Z(K) = 0$.

Proof: Define $S_{in} = \{a \cdot E_{in} \mid a \in D\}$; we want to show that S_{in} is a simple submodule of K_K for $1 \leq i \leq n$. Clearly, S_{in} is closed under addition. Let $k \in K$ and α_{nn} be that element of D that belongs to the n^{th} row and n^{th} column of k . Let $s \in S_{in}$. Then $s = a \cdot E_{in}$ for some $a \in D$. But $s \cdot k$ is the matrix with $a \cdot \alpha_{nn}$ in the i^{th} row and n^{th} column with zero everywhere else. So $s \cdot k \in S_{in}$ and S_{in} is a submodule of K_K . Let $0 \neq L_K$ be a submodule of S_{in} ; we want to show $L = S_{in}$. Because $(S_{in})_K \subseteq K_K$, $L_K \subseteq K_K$. Choose $0 \neq b \in L$. Then $b = a \cdot E_{in}$

for some $a \in D$. Let $b' = a^{-1} \cdot E_{nn}$. Then $b' \in K$ and $E_{in} = b \cdot b' \in L$. Thus, $S_{in} = (b \cdot b')S_{in} \subseteq L$ which implies $S_{in} = L$. Therefore, S_{in} is simple for each $1 \leq i \leq n$.

It is clear that if I is an essential right ideal of K , then $I \cap S_{in} = S_{in}$ because S_{in} is simple for each $1 \leq i \leq n$. This implies $S_{in} \subseteq I$ for each $1 \leq i \leq n$ and hence, $\sum_{i=1}^n S_{in} \subseteq I$.

Observe if c is any element of K , then $c \cdot E_{in}$ is the matrix whose n^{th} column is equal to the i^{th} column of c and is zero everywhere else.

Assume $a \in Z(K)$. Thus $\sum_{i=1}^n S_{in} \subseteq (0:a)$ as shown above. This implies $a \cdot \sum_{i=1}^n S_{in} = 0$ and therefore, $a \cdot E_{in} = 0$ for each $1 \leq i \leq n$. Thus, the matrix a must be equal to zero. Therefore, $Z(K) = 0$.

2.2 LEMMA. $S_K \nabla K_K$.

Proof: It is not hard to show S is a K -module since K is a subring of S . To show $S_K \nabla K_K$, let $0 \neq x \in S$ and prove $K \cap xK \neq 0$. Since $x \neq 0$, there exists an integer j , $1 \leq j \leq n$, such that the j^{th} column of x is not zero. So, $x \cdot E_{jn}$ is the matrix with n^{th} column equal to the j^{th} column of x and zero everywhere else. Thus $x \cdot E_{jn} \neq 0$ and is an element of K . Now, $E_{jn} \in K$ and $x \cdot E_{jn} \in xK$. So $0 \neq x \cdot E_{jn} \in xK \cap K$ and $S_K \nabla K_K$.

2.3 THEOREM. $E(K)$ is a right S -module.

Proof: Since K is a ring and $Z(K) = 0$, by 1.2.6 $E(K)$ is a ring. Since by 2.2, $S_K \nabla K_K$, there exists a function $\phi \in \text{Hom}_K(S, E(K))$ such that $\phi(k) = k$ for each $k \in K$. We will show $E(K)$ is a S -module by first showing multiplication in S is equivalent to multiplication in $E(K)$. Consider the following diagram where θ is the canonical injection map:

$$\begin{array}{ccccc} 0 & \longrightarrow & K_K & \xrightarrow{\theta_K} & S_K \\ & & \downarrow \theta_K & \searrow \phi & \\ & & E(K)_K & & \end{array}$$

We need to show $\phi(ss') = \phi(s)\phi(s')$ for all $s, s' \in S$. Fix $s \in S$. Define $\psi \in \text{Hom}_K(S, E(K))$ by $\psi = \phi \circ \lambda_s$ where $\lambda_s(s') = ss'$ for each $s' \in S$. Define $\psi' \in \text{Hom}_K(S, E(K))$ by $\psi' = \lambda_{\phi(s)} \circ \phi$ where $\lambda_{\phi(s)}(e) = \phi(s) \cdot e$ for all $e \in E(K)$. However, if $k \in K$, $(\psi - \psi')(k) = \psi(k) - \psi'(k) = (\phi \circ \lambda_s)(k) - (\lambda_{\phi(s)} \circ \phi)(k) = \phi(s \cdot k) - \phi(s) \cdot k = \phi(s) \cdot k - \phi(s) \cdot k = 0$. Therefore, $(\psi - \psi')(K) = 0$ which implies, since $(\psi - \psi') \in \text{Hom}_K(S, E(K))$ and $E(K) \nabla K$, that $(\psi - \psi')(S) = 0$. Thus, $\psi(s) = \psi'(s)$ or equivalently if $s \in S$, $\phi(ss') = \phi(s)\phi(s')$. Since s was arbitrary, this is true for each $s \in S$, and ϕ is a ring homomorphism between S and $E(K)$. So, $E(K)$ can be shown to be an S -module by defining $e \mu s = e \cdot \phi(s)$ for all $e \in E(K)$, $s \in S$.

If one considers S_K to actually be a submodule of $E(K)_K$, then to prove $E(K)$ is an S -module one must show $s \cdot s' = s \cdot' s'$

where \cdot is multiplication in S and \cdot' is multiplication in $E(K)$.

2.4 THEOREM. $E(K) = S$.

Proof: By the Wedderburn Theorem, $S = M_n(D)$ is a self-injective right S -module. From the preceding lemmas, we know $E(K)_S \supseteq S_S$. Thus, $E(K)_S = S_S \oplus S'_S$. But any S -module is a K -module since K is a subring of S . Thus, $E(K)_K = S_K \oplus S'_K$. However, since $K \subseteq S$, $K \cap S' = 0$ which implies $S' = 0$, since $E(K) \nabla K$. Therefore, $E(K) = S$.

SUMMARY

In conclusion, we have shown if the right singular ideal of a ring R is zero, then the injective hull of R is a ring whose operations preserve the module structure of the injective hull. Furthermore, this ring is right self-injective and von Neumann.

These results are used to show the injective hull of the $n \times n$ upper triangular matrix ring over a division ring is the full $n \times n$ matrix ring. For an alternate proof of this (nontrivial) result, the reader may consult [1].

BIBLIOGRAPHY

1. Elizabeth Bray, The Injective Envelope of the $n \times n$ Upper Triangular Matrix Ring Over a Field, Master's Thesis, University of North Carolina at Greensboro, Greensboro, N. C., 1970.
2. Carl Faith, Lectures on Injective Modules and Quotient Rings, vol. 49, Springer-Verlag, Berlin, 1967.
3. Mary Gray, A Radical Approach to Algebra, Addison-Wesley, Reading, Mass., 1970.
4. Barbara L. Osofsky, On ring properties of injective hulls, Canad. Math. Bull., vol. 7, no. 3, July, 1964, p. 405.